

FISH FOODS

1997

PRODUCT DATA

PRODUCT DATA			
DIG. ENERGY	1078.07	Kcal/lb	
PROTEIN	32.04	%	
ARGININE	2.01	%	
GLYCINE	1.71	%	
HISTIDINE	0.73	%	
ISOLEUCINE	1.32	%	
LEUCINE	2.34	%	
LYSINE	1.71	%	
METHIONINE	0.44	%	
CYSTINE	0.48	%	
PHENYLALANINE		%	
TYROSINE	0.88	%	
SERINE		%	
THREONINE		%	
TRYPTOPHANE	0.38	%	
VALINE	1.45	%	
FAT	3.13	%	
FIBER	3.92	%	
ASH	7.16	%	
CALCIUM	1.48	%	
PHOSPHORUS	1.04	%	
POTASSIUM	1.30	%	
CHLORINE		%	
MAGNESIUM		%	
SODIUM	0.21	%	
SULFUR	0.30	%	
COPPER	8.06	mg/lb	
IRON	86.05	mg/lb	
MANGANESE	19.05	mg/lb	
SELENIUM	0.15	mg/lb	
ZINC		mg/lb	
BIOTIN	0.15	mg/lb	
CHOLINE	913.22	mg/lb	
FOLIC ACID	1.10	mg/lb	
NIACIN		mg/lb	
PAN. ACID		mg/lb	
VITAMIN B6	6.43	mg/lb	
RIBOFLAVIN	6.19	mg/lb	
THIAMINE	6.14	mg/lb	
VITAMIN B12		mg/lb	
VITAMIN E		mg/lb	
VITAMIN A		KIU/lb	
VITAMIN D3		KIU/lb	
VITAMIN K		mg/lb	
VITAMIN C		mg/lb	

TFP NUTRITION NACOGDOCHES, TEXAS

FLOATING CATFISH FOOD


A ration designed to be fed to catfish 2 months of age and older.

FEATURES

- * 32% Protein
- * 3/16" Pellet
- * Floating ration

BENEFITS

* An economical ration for growing catfish from fingerling size to market.



GUARANTEED ANALYSIS

2.00%	Crude Protein, minimum
3.00%	Crude Fat, minimum
5.00%	Crude Fiber, maximum
0.50%	Phosphorus (P), minimum
3.00 5.00	Crude Fat, minimum

INGREDIENTS

Plant protein products, grain products, processed grain by-products, animal protein products, salt, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, vitamin B12 supplement, riboflavin supplement, niacin supplement, calcium pantothenate, choline chloride, menadione sodium bisulfite complex (source of vitamin K activity), folic acid, pyridoxine hydrochloride, thiamine mononitrate, sodium selenite, biotin, L-ascorbyl-2-polyphosphate (source of vitamin C), manganese sulfate, zinc sulfate, ferrous sulfate, copper sulfate, ethylenediamine dihydroiodide.

FEEDING DIRECTIONS

The amount of feed utilized by the fish depends primarily on water temperature and the size of the fish. However, the rate should not exceed 25 pounds per surface acre of water regardless of the number and weight of the fish. Fish grow best when the water temperature is above 70 degrees F. Typically, the amount of feed taken by the fish will vary directly with the temperature. Use the table below as a guide for feeding channel catfish from fingerlings to market or harvest size. Fish should not be fed more than they will eat in a 24-hour period.

SUGGESTED FEEDING RATES

WATER TEMPERATURE	AMOUNT OF FEED
Above 70 degrees F	3% of estimated total weight of fish
60 to 70 degrees F	2% of estimated total weight of fish
45 to 60 degrees F	1% of estimated total weight of fish
*Below 45 degrees F	Usually no feed

*A small amount of feed may be necessary if low temperature is prolonged. In cold weather, growth will be slow or nonexistent. During extended low water temperatures in the winter, a certain amount of feed is necessary to maintain weight.

If this feeding guide is used, the estimated total weight of all fish in a given pond must be determined each month. A simple procedure can be followed:

- 1. Collect by seine or other method, approximately 100 fish from the pond.
- 2. Count and weigh the fish. Return the fish to the pond as soon as possible.
- 3. Obtain the average weight per fish by dividing the number of fish collected into the pounds of fish collected.
- 4. Obtain estimated total weight by multiplying average weight of a single fish by the total number stacked in the ponds.
- 5. Calculate pounds of feed needed by multiplying the estimated total weight by the percentage of required food indicated in the above table.

Growing fish should be fed 6 days a week and at the same time and place. During cool weather, feed the fish when the highest daily temperature occurs (approximately 3:00 P.M.). Continue feeding until the fish are the desired size. Do not feed to cattle or other ruminants.

CAUTION: Feed should be stored in a well-ventilated, dry area that is protected from rodents and insects. Feed is perishable. Do not feed moldy or insect-infested feed as it may cause illness or death.